



A STUDY OF CHARACTERISTICS OF LIVER ABSCESS IN SURGICAL DEPARTMENT

General Surgery

Dr. Ankitkumar Bagdai	M.S., Assistant Professor, Department of General Surgery, GCS Medical College, Ahmedabad.
Dr. Nehalkumar Sisodiya*	M.S., Assistant Professor, Department of General Surgery, GCS Medical College, Ahmedabad. *Corresponding Author
Dr. Priyanka B. Bagdai	BDS, Department of Dental Science,

ABSTRACT

Introduction: Abscesses of the liver are due to an infection of bacterial, parasitic or fungal origin. Through imaging and serology, their diagnosis was facilitated. The aim of this study was to determine the epidemiological, clinical, diagnostic and evolutionary aspects of these liver abscesses at the Department of general surgery in this Hospital.

Patients and Methods: This was a retrospective and descriptive study, based on records of patients hospitalized for abscess of the liver over a period of 3 years, from May 1, 2015 to June, 2018.

Results: We collected 60 cases of abscesses including 36 cases of confirmed amoebic abscesses and 15 cases of pyogenic abscesses. The mean age of the patients was 43.15 ± 15.12 years. There were 54 men and 6 women. The average hospital stay was 10 days. The most common clinical signs were abdominal pain (100%), fever (75%), hepatomegaly (80%). A leukocytosis greater than 12,000 cell/mm³ was found in 39 patients. The amoebic serology was positive in 71% of cases and negative in 29% of cases. No germ was isolated from the blood culture. On hepatic ultrasound the abscess was unique in 90%. Treatment was exclusively medical in 33 patients (55%). The trend was favorable in 85% of cases.

Conclusion: The incidence of liver abscesses has slightly decreased. It is important to develop diagnostic tools better in the case of pyogenic abscesses.

KEYWORDS

Amoebic Liver Abscess, Pyogenic Abscess,

1. INTRODUCTION

Liver abscesses are suppurations collected in a newly formed cavity within the liver parenchyma due to bacterial, parasitic or fungal infections. Their diagnosis has been facilitated by the progress made over the last decade through the popularization of non-invasive tests such as ultrasounds. Nevertheless, confirmation of the etiology remains difficult due to the limited access to the necessary biological explorations. However, most studies are on amoebic liver abscesses. Improving the living conditions of populations living in endemic areas, particularly increasing access to drinking water and fighting against faecal contamination related diseases should lead to a decline in the prevalence of this condition. More rarely diagnosed in tropical areas, pyogenic abscesses are more common in Europe where their prevalence is between 0.008 and 0.16% in hospitalized patients. This incidence has likely increased according to recent studies. Gram-negative bacilli and anaerobic bacteria are the main culprits in 40 to 60% of cases.

2. PATIENTS AND METHODS :-

This study was conducted at the general surgical department, GCS medical college and researcher hospital, Ahmedabad. It was a retrospective and descriptive study based on records of patients hospitalized at general surgery Unit for liver abscess over an 3 year period, from May 2015 to June 2018. All cases of liver abscesses confirmed during the study period by abdominal ultrasound or CT scan showing an abscessed collection in the liver were included in this study. Patients who had tumor like hepatic masses with high alpha fetoprotein or confirmed by histology were not included. The amoebic etiology was selected in cases of positive serology at a rate of $\geq 1/160$ IU through indirect immunofluorescence, or $\geq 1/320$ IU through hemagglutination, and/or an odorless, anchovy sauce pus, which was sterile when cultured or contained amoebae, obtained using an echo guided puncture or surgical drainage. The pyogenic etiology was selected when the puncture revealed pus and polynuclear are altered with or without germ isolation, and/or a negative amoebic serology. This etiology was also selected in cases of a positive blood culture. Data was obtained from the medical records of hospitalized patients. A standardized data collection form was developed. It included epidemiological aspects (year of occurrence, age, sex, lifestyle, site of occurrence), clinical aspects (time of hospitalization, clinical signs), paraclinical aspects (biological signs: hemogram, VS, disruption, amoebic serology results; ultrasound signs), therapeutic aspects

(medical: type, dose, duration; instrumental; surgical: indication, gesture used) and evolutionary aspects (healing, complication or death). These data was then entered and analyzed on the software.

3. RESULTS

3.1. Epidemiological Aspects: During our study period (2015-2018), we recorded 60 cases of liver abscess among 4219 hospitalized patients. The mean age of the patients was 43.15 ± 15.12 years, with extremes of 19 years and 87 years. There were 54 men and 6 women, among whom 21 patients consumed alcohol, 6 patients had type II diabetes, and 2 patient was living with HIV. Traces of dysenteric syndrome were noted in 15 patients.

3.2. Clinical and Paraclinical Aspects: Patients were hospitalized within 30 days, with extremes of 3 and 180 days. Half of the patients were hospitalized within the first 20 days. Abdominal pain was reported by all patients (100%), located in the right upper quadrant in 42 patients and was generalized to the whole abdomen in 12 patients. 45 patients had a fever, and 48 patients had a clinically detectable hepatomegaly (Table 2). Biologically, the hemogram showed predominantly neutrophilic hyperleucocytosis greater than 12,000 cell/mm³ in 39 patients, and anemia in 42 patients. The chest X-ray showed right-sided pleural effusion in 10 patients, pneumonia in 3 patients, and an elevated hemidiaphragm in 6 patients. According to the ultrasound data, the abscess was unique in 54 cases (90%) and multiple in 6 cases (10%). The right lobe was the preferred seat of these liver abscesses (95%). The abscess size was greater than 10 cm in 20 patients.

Table.1 clinical signs

Clinical signs	Number	Percentage
abdominal pain	60	100
Fever	45	75
Hepatomegaly	48	80
Pleural effusion	10	16.6
Leucocytosis(>12000cell/mm)	39	65
Anemia	42	70

Table.2 disease prevalence in gender

	Number	Percentage
Male	54	90
Female	06	10

3.3. Therapeutic and Evolutionary Aspects:

Treatment was exclusively medical in 33 patients (55%). This medical treatment was associated with surgical drainage in 9 patients (15%) and Ultrasound guided aspirations in 18 patients (30%). Medical treatment with metronidazole-ceftriaxone was the most commonly used (24 cases), followed by metronidazole-ceftriaxone-gentamicin (12 cases) and metronidazole-ampicillin-gentamicin (9 cases). The average hospital stay was 10 days with extremes of 3 days and 30 days. The outcome was favorable in 51 patients, (85% of cases). Complications were noted, such as a pleural rupture in 3 patient, The reactive or purulent nature of the pleural effusions could not be determined because the results of the punctures were unavailable.

DISCUSSION :

From 2015 to 2018, 60 cases of liver abscess were recorded at general surgery department. we observe a decrease of cases over the year's. This decline in incidence may be due to improved standards of individual and collective hygiene, the proliferation of public health facilities and the advent of imidazole derivatives, their popularizations and their easy prescription for any dysenteric syndrome. Of these cases, 36 were confirmed amoebic liver abscesses, and in 15 patients the diagnosis of pyogenic abscess was retained. Amoebic liver abscesses are predominant in the various series reported in Africa in adults [9]. This predominance of cases of amoebic liver abscess can be explained by the high prevalence of intestinal amoebiasis in our regions, which represent the main complication [10]. The average age of 43 years of our patients was comparable to that reported by other authors. In our study, the male sex was predominant, and this finding corroborates data in literature for both amoebic abscess and pyogenic abscess. A history of dysentery was found in 15 patients, i.e. 25% of our patients, which is consistent with the results of Salanke and Laverdant [16] who reported it in 30% and 20.3% of cases respectively. The existence of a history of dysentery or diarrhea should not be excluded in so far as there is no amoebiasis of the liver without colonic amoebiasis. The existence of a lag phase between the two conditions may result in an oversight. Indeed, alcohol intoxication which was noted in 21 patients is a promoting factor of liver abscesses. In India first recourse to traditional healers and self-medication are the main causes of the long delay between the presumed onset of clinical signs and hospitalization. Clinical medicine as classically defined is not always constant. Abdominal pain remains the most frequently encountered sign, although the seat is often atypical, as it was our case. It was reported by all patients in our series and remains the main symptom. This is consistent with the results of other studies. The detection of hepatomegaly is typical but inconstant and was reported in 80% of our patients. This frequency varies according to the studies. Its detection is difficult because of the pain; therefore, its actual rate may be underestimated. The leukocytosis is valuable for orientation but has poor specificity because it does not differentiate etiologically. It was greater than 12,000 Gb/mm³ in 13 patients. This leukocytosis was noted in 92% of Laverdant patients [16] and in 65% of Djossou patients [11]. Amoebic serology, although it is not routinely performed due to its financial inaccessibility and unavailability in public hospitals, It was positive in 36/51 patients i.e. 71%, which is similar to the results of Djossou, where the serology was positive in the same proportions. The detection of serum antibodies confirms amoebic origins. Moreover, the diagnosis of pyogenic abscess was more difficult in our context, based on indirect paraclinical arguments such as a negative amoebic serology and sometimes the appearance of the purulent fluid if a puncture was performed. Detecting the germ using an echo-guided puncture of the abscess remains the determining factor [12]. However, the puncture is not routinely performed, but remains important especially for therapeutic purposes in current practice. Even when it is done, it does not always find the germs in question, but it remains useful in identifying the germ. In Ba's series, the macroscopic appearance of pus was clearly purulent in 12 cases, and the bacteriological study of the puncture fluid identified a bacterium in 5 children [18]. In our series, the blood culture was sterile in all patients which was not great for remembering the etiology of a pyogenic abscess. Blood cultures should be done prior to antibiotic therapy, which is not always the case in current practice, and this reduces the chances of isolating a germ in the case of a pyogenic abscess. In the clinical context, the ultrasound is the exam of choice in the diagnosis and monitoring of liver abscesses due to its easy application, effectiveness and affordable cost. It was performed in all our patients upon admission or during hospitalization, and its sensitivity was 100%. It specified the number, the seat, and the size of the hepatic collections. Thus, we found that in 90% of cases the abscess was

unique. Varying percentages have been reported ranging from 65% to 77% [17] [20] [21]. The right lobe remains the preferred but not exclusive location of liver abscesses [9] [12] [22], which is consistent with our findings where abscesses were present in the right lobe in 95%. The larger volume of the right lobe, the relative prominence of its vascularization, mismatching in the vein carry flows from the superior mesenteric vein and splenic vein, with one flowing electively to the right lobe and the other to the left lobe, are the reasons for this preferential location [19]. The size of the abscess is highly variable and can range from a few millimeters in diameter, to up to 20cm or more [22]. The size of the abscess was greater than 10 cm in 20 patients. Indeed, the size of abscesses could be explained by the delay in consultation, and therefore the care that the patients received. Therapeutically, medical treatment was exclusive in 33 patients i.e. 55% of cases, associated with ultrasound guided aspirations in 30% of cases, and with surgical drainage in 15% of cases. Medical treatment is recommended most often for small abscesses [5] [9]. Metronidazole was the only imidazole used in our study, as in most studies, to target amoebic and anaerobic etiology. Nevertheless, it was combined with Beta-lactamine antibiotic therapy, which in 39 cases was ceftriaxone. This therapeutic approach was performed in the majority of series on liver abscess due to the difficulty of confirmation without completing certain paraclinical exams. The ultrasound guided aspirations is a very commonly used therapeutic method today [19]. The size of the abscesses (abscess \geq 10 cm in diameter), their location in the left lobe, and the persistence of pain are indications of this technique. Surgical drainage is currently the last resort in the treatment of abscesses of the liver and is most often reserved for complications, namely: peritoneal, pleural, pericardial rupture or fistulation of the abscess in a hollow viscus, or on the skin. Surgical drainage was necessary in 9 patients of our series, following a persistence of abscess to medical treatment in 6 cases and a pleural rupture in 3 patient. Overall, liver abscesses have a favorable outcome with proper treatment. 3 case of death was observed in our series. The cases of death described in the literature are secondary to intraperitoneal rupture of the abscess. The favorable evolution of the abscesses of the liver undergoing treatment is clinically assessed by the disappearance of pain and fever in 48 to 72 hours. The regression of hepatomegaly occurs more gradually. C-reactive protein and changes in leukocyte count are always useful for judging short term evolution. As for the ultrasound, it can show the persistence of the abscessed cavity for several months.

5. CONCLUSION

The overall incidence of liver abscess has declined slightly in recent years. Amoebic serology has contributed significantly to the diagnosis of hepatic amoebiasis, hence the need to make it accessible in public hospitals. However, the diagnostic difficulty lies in the confirmation of cases of pyogenic abscess. It is necessary to strengthen and develop other diagnostic methods for these pyogenic abscesses.

REFERENCES

- Service, G., Camengo Police, S.M., Mobima, T., Tekpa, G., Boua Akelelo, N.P. and Koffi, B. (2014) Les abces ambiens du foie dans le service de gastroenterologie a l'hopital de L'Amitie de Bangui: Aspects epidemiologiques, diagnostiques, therapeutiques etevolutifs. *Journal Africain d'Imagerie Medicale*, 6, 4-9.
- Traore, A., Togo, A., Kante, L., Demebele, B.T., Diakite, I., Coulibaly, Y., et al. (2014) Abces ambien du foie en chirurgie generale du CHU Gabriel Toure, Bamako. *Mali Medical*, 29, 5-9.
- Kodjoh, N. and Hountondji, A. (1991) Les abces ambiens du foie a propos de 42 observations colligees dans les services medicaux du CHNU de Cotonou. *Medecine d'Afrique Noire*, 38, 185-190.
- Pelissier, R. (1990) La ponction echoguidee: Traitement moderne des abces du foie. *Annales De Radiologie*, 33, 277-279.
- Greenstein, A.J. and Sachar, D.B. (1988) Pyogenic and Amoebic Abscess of the Liver. *Seminars in Liver Disease*. *Liver Disease Journal*, 8, 210-217.
- Charigneau Derodde, C. and Sylvain, C. (2004) Abces bacteriens du foie. *Gastroenterologie Clinique Et Biologique*, 28, 470-476. [https://doi.org/10.1016/S0399-8320\(04\)94961-5](https://doi.org/10.1016/S0399-8320(04)94961-5)
- Lee, K.T., Sheen, P.C., Chem, J.S. and Ker, C.G. (1991) Pyogenic Liver Abscess: Multivariate Analysis of Risk Factors. *World Journal of Surgery*, 15, 372-377. <https://doi.org/10.1007/BF01658732>
- Dieng, M., Diop, B., Konate, I., Ka, O., Dia, A. and Toure, C.T. (2007) Traitement des abces du foie: L'experience d'un service de chirurgie generale. *Medecine d'Afrique Noire*, 54, 513-519.
- Nozais, J.P., Datry, A. and Danis, M. (1996) *Traite de parasitologie*. Editions Pradel, Paris, 249-251.
- Djossou, F., Malvy, D., Tamboura, M., Beylot, J., Lamouliatte, H., Longy-Boursier, M., et al. (2003) Abces ambien du foie. Analyse de 20 observations et proposition d'un algorithme therapeutique. *La Revue de Medecine Interne*, 24, 97-106. [https://doi.org/10.1016/S0248-8663\(02\)00015-2](https://doi.org/10.1016/S0248-8663(02)00015-2)
- Ibara, J.R., Ollandzobo Ikobo, L.C., Atipo Ibara, B.I. and Itoua Ngaporo, A. (2000) Abces du foie a germes pyogenes aspects cliniques, morphologiques et etologiques a propos de 38 cas. *Medecine d'Afrique Noire*, 47, 92-96.
- Nguema-mve, R. and Graesslin, O. (1997) Traitement des abces ambiens du foie par le drainage percutane. *Medecine d'Afrique Noire*, 44, 36-39.
- Lebeau, R., Diane, B., Ynon, S., Koffi, E., Kassi, B.A. and Kouassi, J.C. (2014) Abces ambien du foie au CHU de Bouake: Etude diagnostique et therapeutique a propos de 46

- cas.Mali Medical , 21, 10-14. V. M. P. C. Diallo et al. DOI: 10.4236/aid.2018.81004 31
Advances in Infectious Diseases
- [14] Sankale, M., Diop, B. and Diouf, S. (1970) Reflexions sur l'amibiase hepaticque a Dakar. A propos de 223 cas observes en 10 ans. *Medecine d'Afrique Noire* , 17, 467-477.
- [15] Laverdant, C., Denece, J.M., Roue, R., et al. (1984) L'amibiase hepaticque: etude de 152 observations. *Gastroenterologie Clinique et Biologique* . 8, 634-844.
- [16] Oudou, N., Elie, C., Ndjitoyap, N., Vincent, N. and Charlotte, N. (1999) Abces amibiens du foie a Yaounde. *Cahier Sante* , 9, 119-122.
- [17] Ba, I., Sagna, A., Thiongane, A., et al. (2015) Abces du foie chez l'enfant au Senegal. *Revue Cames sante* , 23, 17-22.
- [18] Molinie, C. and Mennelier, D. (1997) Amibiase hepaticque. *Encyclopedie Medico-Chirurgicale*.
- [19] Weinke, T., Grobusch, M.P. and Guthoft, W. (2002) Amoebic Liver Abscess-Rare Need for Percutaneous Treatment Modalities. *European Journal of Medical Research*
- [20] Lodhi, S., Sarwari, A.R., Muzammil, M., Salam, A. and Smego, R.A. (2004) Features Distinguishing Amoebic from Pyogenic Liver Abscess: A Review of 577 Adult Cases. *Tropical Medicine and International Health* .
- [21] Le Bras, Y., Gervez, F., Abraham, E., Toussaint, T. and Verhaegen, F. (1991) Interet de la ponction evacuatrice echoguidee dans le traitement des abces amibiens du foie. *Journal de Radiologie* , 72, 43-47.
- [22] Chuah, S.K., Chang-Chien, C.S., et al. (1992) The Prognostic Factors of Severe Amebic Liver Abscess: A Retrospective Study of 125 Cases. *The American Journal of Tropical Medicine and Hygiene* , 46,
- [23] Lee, C.K., Yamazaki, O., Hamba, H., et al. (1996) Analysis of 69 Patients with Amebic Liver Abscess. *Journal of Gastroenterology* , 3