



ORIGINAL RESEARCH PAPER

ENT

PSEUDOMONAS AERUGINOSA; MICROBIOLOGICAL PROFILE AND ANTIMICROBIAL SENSITIVITY PATTERN IN CHRONIC SUPPURATIVE OTITIS MEDIA

KEY WORDS:

Dr Prathap R

Associate Professor Karuna Medical College, Chittur Vilayodi ,678103

Dr Asim P A*

Assistant Professor Karuna Medical College, Chittur Vilayodi ,678103
*Corresponding Auhtor

Dr Midun Mohandas

Senior Resident Karuna Medical College, Chittur Vilayodi ,678103

ABSTRACT

INTRODUCTION: Chronic Suppurative Otitis Media (CSOM) is a potentially serious disease because of its complications. The incidence of CSOM is increasing in the developing countries because of the poor hygiene practices and lack of health education.

Pseudomonas aeruginosa is a Gram-negative bacterium belonging to the family Pseudomonadaceae that is able to survive in a wide range of environments.

AIM: To determine the microbiological profile and antimicrobial sensitivity pattern of *Pseudomonas aeruginosa* from discharge in CSOM.

MATERIAL AND METHOD: This study was conducted over a time period of 6 months. All bacterial isolates were identified by conventional microbiological methods. Antimicrobial sensitivity testing of the isolates was done by Kirby-Bauer disc diffusion method according to Clinical and Laboratory Standards Institute (CLSI) guidelines.

RESULT: *Pseudomonas aeruginosa* (58.6 %) was the most prevalent bacteria isolated, followed by staphylococcus aureus (19.57%). Most sensitive antibiotic against *Pseudomonas aeruginosa* was Piperacillin – Tazobactam (76.5%) followed by Meropenam (65.8%) and cephalosporins such as cephalexin, cefuroxime. Amoxyllin-clavulanic acid shows complete resistance towards *Pseudomonas aeruginosa*.

CONCLUSION: It is important to evaluate bacteriologic profile and antimicrobial susceptibility pattern periodically for efficacious treatment in CSOM patients

INTRODUCTION

Among the various organisms isolated, *Pseudomonas aeruginosa* has been particularly blamed for deep seated and progressive destruction of middle ear and mastoid structures by releasing its toxins [1]. *Pseudomonas aeruginosa* is a Gram negative bacterium belonging to the family Pseudomonadaceae that is able to survive in a wide range of environments [2]. The mechanism for antibiotic resistance can be classified into intrinsic, acquired and adaptive resistance. The intrinsic resistance of *P. aeruginosa* includes low outer membrane permeability, expression of efflux pumps that expel antibiotics out of the cell and the production of antibiotic inactivating enzymes. The acquired resistance of *P. aeruginosa* can be achieved by horizontal transfer of resistance genes [3]. The adaptive resistance of *P. aeruginosa* involves formation of biofilm in the lungs of infected patients where the biofilm serves as a diffusion barrier to limit antibiotic access to the bacterial cells [4]. Development of complications depends on various reasons like high virulence of organism, poor resistance of patients, inadequate antibiotic treatment of acute middle ear and mastoid infection, presence of chronic systemic diseases and resistance of organisms to antibiotics which are becoming common on these days [5].

MATERIALS AND METHODS

A prospective observational study was conducted in the department of ENT at Karuna Medical College Hospital, Vilayodi, Chittur, Kerala, over a period of 6 months from October 2018 to March 2019. This study was conducted after getting approval from the Institutional Human Ethical Committee (IHEC/09/2018) of KMCH. The patients attending in the ENT department with active purulent ear discharge and clinically diagnosed as squamosal, mucosal type of CSOM and malignant otitis externa was included in this study. Patients having malignancy of middle ear, previous history of surgery and Patient who already received treatment within 2 week for the same complaints of otitis media and malignant otitis externa were excluded from this study. Each of these

patients were subjected to detailed clinical history regarding to name, age, sex, laterality and symptoms like ear discharge, hearing loss, vertigo, head ache , fever , nausea & vomiting, and facial weakness. Thorough ENT examination was done for each case by otoscopy and microscopic examinations. Pure tone audiometry (PTA) carried out for relevant cases. Two sterile cotton swabs were used to collect ear discharge from CSOM patients. One swab was used for performing gram stain and second one for culture. The culture was done on nutrient agar, blood agar, MacConkey agar and SDA. The organisms were identified by culture characteristics, morphology and conventional biochemical tests. Antimicrobial susceptibility testing for *pseudomonas aeruginosa* was performed on Mueller Hinton agar plate using Kirby-Bauer disc diffusion method. Collected data was entered in excel sheet for further evaluation and the results was presented in percentage. The main aim of the study was to determine the antimicrobial sensitivity pattern of *Pseudomonas aeruginosa* in CSOM patients.

RESULTS

In the present study of '*pseudomonas aeruginosa*; micro bio logical profile and antimicrobial sensitivity pattern in chronic suppurative otitis media', conducted from October 2018 to March 2019; 123 patients fulfilling the inclusion criteria were randomly selected and analyzed. A total of 246 swabs were collected from 123 cases and sent for gram staining and culture sensitivity. Culture of 123 swabs showed 25 (20.3%) were sterile while 98 swabs yielded growth of organisms. Out of 123 swabs, 3 (2.45%) yielded polymicrobial organisms, while 92 (74.80%) yielded mono microbial flora and three swabs shows fungal growth (2.45%). To detect the presence and identify the general type of bacteria or sometimes fungi (microbes) in a sample taken from the site of a suspected infection; to generally classify bacteria grown in culture so that further identification tests can be performed and appropriate treatment given. Polymicrobial isolation was associated with 6 organisms i.e. one combination of *Pseudomonas spp.* and *Klebsiella species*. Another two

combinations include Klebsiella spp. with coagulase negative staphylococcus aureus and streptococcus species. So out of total microbes isolated (n=98), 73 were gram -ve bacteria and 25 were gram +ve organisms . Among the mono microbial isolates, Pseudomonas aeruginosa was the commonest offender seen in 54 (58.6%) swabs, followed by Staphylococcus aureus in 18 (19.57%) swabs. The most predominant organism causing CSOM among aerobic bacteria was Pseudomonas aeruginosa, followed by Staphylococcus aureus .

Most of the Pseudomonas species were sensitive to piperacillin-tazobactam (76.5%). Meropenam were the second most effective group, showing 65.8% sensitivity followed by tobramycin (64.7%), and ceftazidime-sulbactam (56.7%) respectively. Cephalosporins such as cephalexin, cefuroxime and amoxicillin-clavulanic acid shows 100 % resistance to P.aeruginosa. Thus the most sensitive antibiotics to P. aeruginosa were piperacillin-tazobactam and the least effective were cephalexin, cefuroxime and amoxicillin-clavulanic acid.

DISCUSSION

Analysis of bacteriology culture results from our study found pure culture (Monomicrobial) to be more common (74.80%) and this observation is supported by other investigators [6-9]. In the present study, out of total microbes isolated (n=98), 73 were gram -ve bacteria and 25 were gram +ve organisms. And which is in concordance with findings of other study which shows out of 90 culture smears, 59 (65.55%) were Gram negative and 31 (34.45%) were Gram positive [10].

Our study reveals that among the mono microbial isolates, Pseudomonas aeruginosa was the commonest offender seen in 54 (58.6%) swabs, followed by Staphylococcus aureus in 18 (19.57%) swabs. This result was comparable with study conducted by Bairy I et al which shows Pseudomonas aeruginosa (33.9%) as the commonest organism [11]. Similarly, Pseudomonas was also isolated as the predominant organism by Maji PK et al (63.8%) [12], Kumar S et al (45.9%) [13] and Indudharan R et al (27.2%) [14].

In our study Citrobacter species (1.08%) was found as the least isolates among monomicrobial organisms which is comparable with study conducted by Kumar R et al [15].

Pseudomonas aeruginosa showed higher resistance to cephalosporins such as cephalexin, cefuroxime and amoxicillin-clavulanic acid and it is supported by various studies like Ajay K et al [16] and higher sensitive to piperacillin-tazobactam (76.5%) followed by meropenam showing 65.8% sensitivity followed by tobramycin (64.7%), and ceftazidime-sulbactam (56.7%) respectively and it was comparable with study conducted by Mansoor T et al [17].

CONCLUSION

CSOM is one of the commonest chronic infectious diseases in developing countries. Factors playing role in the pathogenesis of CSOM are not fully comprehensible and underscore the urgent need for research in the area of development of novel and effective therapeutic strategies. Selection of antimicrobial agents must factor in the problem of drug resistance among infecting organisms. Hence, it becomes imperative on part of the health care facilities to undertake periodic evaluation of antimicrobial susceptibility profiles of the microbial pathogens for designing effective empiric treatment protocols and prevent potential risk of unforeseen complications. We conclude that the Pseudomonas aeruginosa was moderately sensitive to Piperacillin-Tazobactam, Meropenam, Tobramycin and Ceftazidime-sulbactam respectively. Resistance towards quinolones group of antibiotics increases when comparing with other studies. Regular antimicrobial sensitivity studies should be conducted as the microbial growth pattern changes with time

because of indiscriminate and haphazard use of antibiotics. This study is more relevant in present scenario, because the knowledge of antimicrobial sensitivity pattern of microorganisms of CSOM will guide clinician in prescribing an empirical regimen so that a more targeted therapy can be provided to CSOM patients.

REFERENCES:

1. Dr. Nawaz Umar, Bilal Ahmad Mir. Prevalence and antibiogram of Pseudomonas aeruginosa isolated from otitis media in KBN hospital, Gulbarga. Sch J App Med Sci 2015;3(7A):2502-2505.
2. Silby MW, Winstanley C, Godfrey SA, Levy SB, Jackson RW. Pseudomonas genomes: diverse and adaptable. FEMS Microbiol Rev 2011;35:652-680.
3. Breidenstein EB, de la Fuente-Nunez C, Hancock RE. Pseudomonas aeruginosa: all roads lead to resistance. Trends Microbiol 2011;19:419-426.
4. Drenkard E. Antimicrobial resistance of Pseudomonas aeruginosa biofilms. Microbes Infect 2003;5:1213-1219.
5. Swapnil R. Pawar, Yogesh Shukla. A clinical study on complications of chronic suppurative otitis media and level of awareness in patients admitted at tertiary care hospital in central India. Int J Community Med Public Health 2015;2(3):223-227.
6. Shyamala R, Reddy PS. The study of bacteriological agents of chronic suppurative otitis media--aerobic culture and evaluation. J Microbiol Biotechnol Res 2017;2:152-62.
7. Poorey VK, Iyer A; Study of bacterial flora in CSOM and its clinical significance. Indian J Otolaryngol Head Neck Surg 2002;54(2):91-95.
8. Prakash M, Lakshmi K, Anuradha S, Swathi GN. Bacteriological profile and their antibiotic susceptibility pattern of cases of Chronic Suppurative Otitis Media. Asian J Pharma Clin Res 2013;6(3):210-212.
9. Mansoor T, Musani MA, Khalid G, Kamal M. Pseudomonas aeruginosa in chronic suppurative otitis media: Sensitivity spectrum against various antibiotics in Karachi. J Ayub Med Coll Abbottabad 2009;21(2):120-123.
10. Wadile Rahul Gopichand, Bhathe Viraj Madhusudan and Kalsheetti Varsha Tukaram. Bacteriological Profile of Chronic Suppurative Otitis Media. Int J Curr Microbiol App Sci 2015;4(6):41-47.
11. Bairy I, Pradhan D, Yenigalla BM. Microbiology of chronic suppurative otitis media. Int J Otol 2007;13:21-4.
12. Maji PK, Chatterjee TK, Chatterjee S, et al. The investigation of bacteriology of chronic suppurative otitis media in patients attending a tertiary care hospital with special emphasis on seasonal variation. Ind J Otolaryngol Head Neck Surg 2007;59(2):128-31.
13. Kumar S, Sharma R, Saxena AK, et al. A study of bacterial flora and sensitivity to antibiotics in cases of CSOM TTD in western UP. Ind J Otol 2016;14:20-4.
14. Indudharan R, Haq JA, Aiyar S. Antibiotics in chronic suppurative otitis media: a bacteriologic study. Ann Otol Rhinol Laryngol 1999;108(5):440-5.
15. Rakesh kumar, P. srivastava, M. sharma, S. Rishi, PS. Nirwan, K. Hemwani, SS. Dahiya. Isolation of antimicrobial sensitivity profile of bacterial agents in chronic suppurative otitis media patients at NIMS hospital, Jaipur. IJPBS 2013;3(4):265-269.
16. Kumar Ajay, Kumari Namrata, Kumar Shailesh, Kumar Rakesh, Xess Anima, Shahi SK. Prevalence and antimicrobial sensitivity pattern of Pseudomonas aeruginosa from chronic suppurative otitis media (CSOM) cases in a tertiary care hospital of Bihar. J Evol Med Dent Sci 2017;8:70.
17. Mansoor T, Musani MA, Khalid G, Kamal M. Pseudomonas aeruginosa in chronic suppurative otitis media: sensitivity spectrum against various antibiotics in Karachi. J Ayub Med Coll Aotabbad 2009;21(2):120-23.